

HATCHGARD®-3 Newcastle - Bronchitis Vaccine B1 Type, B1 Strain, Mass and Conn Types Live Virus

Product Information

HATCHGARD®-3 combines a highly immunogenic isolate of B1 strain Newcastle disease (ND) virus with the well-established infectious bronchitis (IB) protection provided by the Intervet mild Massachusetts and Connecticut types of IB virus. It is a frozen vaccine approved for coarse spray vaccination of chickens one day of age for protection against Newcastle disease and infectious bronchitis (Massachusetts and Connecticut types).



HATCHGARD-3 contains:

- B1 Type, B1 Strain Newcastle Disease virus.
- Massachusetts type of infectious bronchitis virus.
- Connecticut type of infectious bronchitis virus.

Product Features:

- Safe to use for hatchery application.
- Frozen for consistent, dependable titer every serial.
- Mild reaction.
- Highly immunogenic protection against ND and a broad spectrum protection against IB.

Indications for Use:

HatchGard-3® is indicated for the vaccination of chickens at one day of age for the prevention of Newcastle disease and infectious bronchitis (Massachusetts and Connecticut types).

Dosage and Administration:

HatchGard-3 is approved for use via coarse spray at one day of age. Thaw the vaccine immediately before use. Care should be taken to mix all of the vaccine from the ampule into the diluent for maximum efficacy.

Vaccination:

The best protection is obtained when the vaccinated chickens have time to develop immunity prior to exposure to Newcastle disease or infectious bronchitis virus. Good management practices are recommended to reduce the exposure to these viruses for at least 3 weeks following vaccination. Chickens should be revaccinated with a field boost for optimum protection.

Cautions:

1. Refer to package insert for specific directions.
2. Vaccine should be stored in liquid nitrogen at a temperature below -150°C.
3. Gloves and visor should be worn when handling liquid nitrogen.
4. *Once thawed, this product should not be refrozen.*
5. Vaccinate only healthy birds.
6. Efforts should be taken to reduce stress conditions at the time of vaccination and during the reaction period.
7. Do not dilute the vaccine or otherwise stretch the dosage of the vaccine.
8. Use entire contents of the vial within 2 hours. Unused vaccine should be discarded into disinfectant or burned.
9. Do not vaccinate chickens within 21 days of slaughter.
10. **WARNING:** *Newcastle virus occasionally causes conjunctivitis in humans. Avoid any contact of the vaccine with eyes.*

How Supplied

1 x 10,000 dose ampules


Product Code

NB-226-10

Tab



Cane



Ampule

