

| PRODUCT | <i>Presentations</i> | <i>Dose</i> | <i>Booster Dose</i> | <i>Route of Administration</i> | <i>Clostridium chauvoei (Blackleg)</i> | <i>C. septicum (Malignant edema)</i> | <i>C. novyi Type B (Black disease)</i> | <i>C. sordellii</i> | <i>C. perfringens Types C & D (Enterotoxemia)</i> | <i>Moraxella bovis (Pinkeye*)</i> | <i>Moraxella bovoculi</i> |
|--|----------------------|-------------|-----------------------|--------------------------------|--|--------------------------------------|--|---------------------|---|-----------------------------------|---------------------------|
| BOVILIS® PILIGUARD® PINKEYE+7 | 10-dose 50-dose | 5 mL | 5 mL, 3-4 weeks later | SQ† | X | X | X | X | X | X | |
| BOVILIS® PILIGUARD® PINKEYE | 10-dose 50-dose | 2 mL | | SQ† | | | | | | X | |
| BOVILIS® 20/20 VISION® 7 WITH SPUR® | 10-dose 50-dose | 2 mL | 2 mL, 3-4 weeks later | SQ† | X | X | X | X | X | X | |
| MORAXELLA BOVOCULI BACTERIN | 10-dose 50-dose | 2 mL | 2 mL, 21 days later | SQ† | | | | | | | X |

*Infectious bovine keratoconjunctivitis

†Subcutaneous injection