

All the expertise and support of Merck Animal Health plus the benefits of tulathromycin.



UIN	Product Description	Product Size	Distributor UOM
352606	AROVYN™ 1 x 50 mL (120 bottles per case)	50mL	BTL
382608	AROVYN™ 1 x 100 mL (40 bottles per case)	100mL	BTL
355595	AROVYN™ 1 x 250 mL (24 bottles per case)	250mL	BTL
359543	AROVYN™ 1 x 500 mL (12 bottles per case)	500mL	BTL

Available for order on May 2nd, with estimated shipping beginning the week of May 9th.

→ AROVYNforBRD.com

DOSAGE & ADMINISTRATION: Inject subcutaneously as a single dose in the neck at a dosage of 2.5 mg/kg (1.1 mL/100 lb) body weight (BW). Do not inject more than 10 mL per injection site.

IMPORTANT SAFETY INFORMATION: AROVYN has a pre-slaughter withdrawal time of 18 days in cattle. Do not use in female dairy cattle 20 months of age or older. Do not use in animals known to be hypersensitive to the product. See Full Prescribing Information.

MAHCattle.com · 800-521-5767 · © 2022 Merck & Co., Inc., Rahway, NJ, USA and its affiliates. All rights reserved. US-ARV-220400003

