

MILDVAC[®] - MASS+ARK[®]


Bronchitis Vaccine

(Mild Massachusetts and Arkansas Types, Live Virus)

For the vaccination of healthy chickens by coarse spray at 1 day of age or older as an aid in the prevention of infections due to the Massachusetts and Arkansas type infectious bronchitis (IB) viruses.

ADVANTAGES:

- Mild reactions for minimal stress
- Reduces production losses and performance setbacks
- 2 strains combined provide excellent cross protection; saves time in mixing



Mildvac[®]-Mass+Ark is a lyophilized vaccine. It contains the mild Massachusetts and Arkansas type IB viruses.

10 x 20,000 doses

MILDVAC[®] - MASS+ARK[®]

Bronchitis Vaccine

(Mild Massachusetts and Arkansas Types, Live Virus)

For Animal Use Only.

Indications for Use

Mildvac[®]-Mass+Ark is indicated for the vaccination of healthy chickens by coarse spray at 1 day of age or older as an aid in the prevention of infections due to the Massachusetts and Arkansas types of infectious bronchitis viruses.

Dosage and Administration

The time of administration should be determined by the degree of infectious bronchitis exposure, type of chicken being vaccinated, and the current management procedures. Mildvac-Mass+Ark is approved for use via coarse spray in chickens at 1 day of age or older.

Vaccination

The best protection is obtained when the vaccinated chickens have time to develop immunity prior to exposure to infectious bronchitis virus. Good management practices are recommended to reduce the exposure to this virus for at least 3 weeks following vaccination.

Caution

1. Refer to the package insert for specific directions.
2. Vaccinate only healthy birds. Although disease may not be evident, concurrent disease conditions may cause complications or reduce immunity.
3. All susceptible birds on the same premises should be vaccinated at the same time.
4. Efforts should be taken to reduce stress conditions at the time of vaccination and during the reaction period.
5. Do not spill or splash the vaccine.
6. Do not dilute the vaccine or otherwise stretch the dosage.
7. Use entire contents when first opened.
8. Do not vaccinate within 21 days of slaughter.
9. Burn containers and all unused contents.
10. This vaccine contains gentamicin as a preservative.
11. Avoid exposure of this vaccine to sunlight.
12. Do not use less than one dose per bird.
13. The revaccination of laying hens with live bronchitis vaccine may be detrimental to the flock and cannot be generally recommended.

Notice

This vaccine has undergone rigid potency, safety and purity tests, and meets Merck Animal Health and USDA requirements. It is designed to stimulate effective immunity when used as directed, but the user must be advised that the response to the product depends upon many factors, including, but not limited to, conditions of storage and handling by the user, administration of the vaccine, health and responsiveness of individual chickens, and the degree of field exposure. Therefore, directions should be followed carefully! This product is not hazardous when used according to directions supplied. A safety data sheet (SDS) is available upon request. This and any other consumer information can be obtained by calling Merck Animal Health Customer Service at 1-800-211-3573.

Records

Keep a record of vaccine, quantity, serial number, expiration date, and place of purchase; the date and time of vaccination; the number, age, breed and locations of chickens; name of operators performing the vaccination and any observed reactions.

STORE VACCINE BETWEEN 2 AND 7°C (35 and 45°F).

Contact our sales or technical services representatives to help design a custom vaccination program.