

panacur[®]C (fenbendazole)

The Trusted Dog Dewormer

Indications

For the treatment and control of:



Roundworms

(*Toxocara canis*,
Toxascaris leonina)



Hookworms

(*Ancylostoma caninum*,
Uncinaria stenocephala)



Whipworms

(*Trichuris vulpis*)



Tapeworms

(*Taenia pisiformis*)



- Available in 3 sizes of convenient pre-measured packets by weight (1gm, 2gm, 4gm)
- Safe for use in puppies 6 weeks or older.
Do not deworm a dog or a puppy that is sick
- Repeat the daily dose for 3 days in a row

Approximately 1% of dogs had vomiting associated with the use of this product. PANACUR[®] C (fenbendazole) Canine Dewormer has no known drug interactions and is an ideal complement to monthly heartworm prevention products, which may not protect against all four major types of intestinal worms. See Package Insert on back for full information including side effects, precautions, warnings and contraindications.

Copyright © 2022 Merck & Co., Inc., Rahway, NJ, USA and its affiliates.
All rights reserved. US-PAN-220500001



PANACUR® C (fenbendazole) Intervet/Merck Animal Health

Product Information

Canine Dewormer

Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

Drug Facts

Active ingredients (in each dosage unit):

Fenbendazole Granules 22.2% (222 mg/g)

Purpose: Dewormer for Dogs Only

Uses: For the treatment and control of Roundworms (*Toxocara canis*, *Toxascaris leonina*), Hookworms (*Ancylostoma caninum*, *Uncinaria stenocephala*), Whipworms (*Trichuris vulpis*), and Tapeworm (*Taenia pisiformis*).

Human Warning: Keep this and all medication out of the reach of children.

Adverse Reactions: In US clinical studies 3 of 240 dogs (about 1% of the treated dogs) had vomiting associated with use of the product.

When using this product: Panacur® C is safe for use in puppies 6 weeks or older and adult dogs, including pregnant bitches. Do not deworm a dog or puppy that is sick. Consult your veterinarian for diagnosis of the illness.

Directions:

The daily dose for Panacur® C is 50 mg/kg (22.7 mg/lb) of body weight. Please refer to the following dosing table for help in finding the right dose for your dog.

Dosing Table

Dog Weight	Packet Size*
10 pounds	1 gram
20 pounds	2 gram
30 pounds	1 gram + 2 gram
40 pounds	4 gram
50 pounds	1 gram + 4 gram
60 pounds	2 gram + 4 gram
80 pounds	Two 4 gram
Over 80 pounds	Use combinations to obtain recommended daily dose

*Packet size is the **daily dose**. The dog must be treated with this dose for 3 days in a row.

- You should weigh your dog to make sure you are using the right size and number of packets. If your dog's weight is in-between the suggested dosing sizes, it is safe to use the next higher size. For example, a 15-pound dog should be treated with the 2-gram packet.
- Give Panacur® C to your dog by mixing the daily dose with a small amount of the usual food. Make sure your dog eats all of the medicated food. You may need to moisten dry dog food to aid mixing.
- Repeat the daily dose for three days in a row.

Other Information:

Diagnosis of Parasites: Specific diagnoses require laboratory testing. Consult your veterinarian for help in the diagnosis, treatment, and control of parasitism. Dogs can become infected with several different kinds of tapeworms. Panacur® C only kills *Taenia* species of tapeworms. If you continue to see tapeworm segments in your dog's stool after treatment with Panacur® C, consult your veterinarian.

Recommended Deworming Schedule: Deworming schedules may vary depending on the climate where you live and the activity of your dog. The following schedule should be used as general guidance. Newly weaned pups (6 to 8 weeks of age) should be dewormed at 6, 8, 10, and 12 weeks of age. Treat the dam at the same time as the puppies. Dogs over six months of age should be dewormed at least twice each year. Each deworming requires 3 daily treatments. (See Directions)

Storage: Store at controlled room temperature 68-77°F (20-25°C).

Questions? Comments?

To report a suspected adverse reaction or to obtain product information including material safety data sheets (MSDS), call 1-800-441-8272.

Manufactured by: Intervet GesmbH, Vienna, Austria

Made in Austria

Distributed by: **Intervet Inc d/b/a Merck Animal Health**, Madison, NJ 07940

Fenbendazole (active ingred.) made in China. Formulated in Austria.

Approved by FDA under NADA # 121-473

Net Weight:	Package Contents:	
3 grams	Three 1-gram packets Each Packet Treats 10 lbs.	399072 R4 Rev. 03/21 446500-C
6 grams	Three 2-gram packets Each Packet Treats 20 lbs.	349623 R4 Rev. 03/21 446600-C
12 grams	Three 4-gram packets Each Packet Treats 40 lbs.	366041 R4 03/21 446700-C

CPN: 1047431.2

