

Porcilis® Ileitis

LAWSONIA INTRACELLULARIS BACTERIN



For use in healthy swine as an aid in the control of ileitis caused by *Lawsonia intracellularis*, an aid in the reduction of colonization by *Lawsonia* and an aid in the reduction of duration of fecal shedding. Duration of immunity for at least 20 weeks has been demonstrated.

Advantages:

- Porcilis Ileitis offers long-lasting protection for ileitis control
- Industry-leading 20-week (5 months) duration of immunity (DOI)
 - With a 20-week DOI, Porcilis Ileitis provides 13 more weeks of protection than the water-administered vaccine on the market
- One-dose, ready-to-use injectable vaccine
 - As the only injectable vaccine available for ileitis, Porcilis Ileitis allows for vaccination of pigs at 3 weeks of age or older — an age when they're easier to handle
- Allows pigs to remain on feed and water antibiotics during one of the most challenging times — post-weaning
- No concerns about water quality or water intake
- Microsol Diluvac Forte® adjuvant provides optimal immune response



Code: 126464 100mL
50 doses - 2mL

Code: 144648 500mL
250 doses - 2mL

DRIVEN BY



PREVENTION™



Porcilis® Ileitis

LAWSONIA INTRACELLULARIS BACTERIN



For use in healthy swine as an aid in the prevention of viremia, aid in the reduction of virus shedding and aid in the reduction of lymphoid infection cause by PCV2. Efficacy for at least 20 weeks following completion of vaccination has been demonstrated.

ADMINISTRATION AND DOSAGE:

Shake well. Warm the vaccine to room temperature immediately prior to administration. Using aseptic technique, inject 2 mL once intramuscularly (IM) into pigs at 3 weeks of age or older.

CAUTIONS:

Store in the dark at 2° to 7°C (35° to 45°F). Do not freeze. Use entire contents when first opened; do not save partial contents. Burn the container and all unused product. Use only in healthy swine. Do not vaccinate within 21 days before slaughter. A significant number of pigs may experience transient swelling at the site of injection. The swelling should resolve without treatment in less than 20 days. Immediate systemic reactions have been observed in animals administered this product. Minimize pig handling and ensure proper ventilation prior to starting vaccination. If an allergic reaction occurs, treat with epinephrine. Caution should be used to avoid injecting your own fingers, hands, or other body parts. Accidental injection can cause serious local reactions. Contact a physician immediately.

PRESERVATIVE: Gentamicin and thimerosal.

FOR ANIMAL USE ONLY

READ AND FOLLOW LABEL DIRECTIONS

U.S. Veterinary License No. 165A

FOR VETERINARY USE ONLY

Intervet Inc. d/b/a Merck Animal Health
Omaha, Nebraska 68103 USA
1 800 521-5767