

# Live Free

Keep Your Dog Safe by  
controlling Intestinal Parasites

**safe-guard**<sup>®</sup>  
(fenbendazole)  
**Canine Dewormer**



The dogs we love are more than our companions and family, they're our partner in an outdoor lifestyle. SAFE-GUARD<sup>®</sup> (fenbendazole) helps ensure there's no need to worry about dangerous intestinal parasites getting in the way of enjoying outdoor adventures together.



**Treats More  
Worms**

SAFE-GUARD treats 6 different intestinal worm types common to outdoor dogs with only one compound, fenbendazole. So you can treat more with less, minimizing your dog's chemical exposure.<sup>1</sup>



**Proven  
Safety**

SAFE-GUARD Canine Dewormer is safe even for puppies and pregnant dogs and has no known drug interactions.<sup>1</sup>



**Ease of  
Application**

Dogs readily accept SAFE-GUARD Canine Granules. When mixed in food, dogs usually eat it without fuss.<sup>1</sup>



<sup>1</sup>SAFE-GUARD (fenbendazole) product label







**IMPORTANT SAFETY INFORMATION:** Consult your veterinarian for assistance in the diagnosis, treatment and control of parasitism.



# Color Your Way to the Perfect Dose

Getting the perfect dose with SAFE-GUARD® is easy. Just find your dog's weight range below and select the indicated boxes.



DOG WEIGHT	PACKET SIZE	PACKAGE
10 LBS	1 gram	
11-20 LBS	2 gram	
21-30 LBS	1 gram + 2 gram	
31-40 LBS	4 gram	
41-50 LBS	1 gram + 4 gram	
51-60 LBS	2 gram + 4 gram	
61-80 LBS	4 gram + 4 gram	
80+ LBS		Use combination to obtain recommended daily dose

## Application Instructions

1. Weigh your dog to make sure you are using the correct dosage.
2. Follow the box directions for daily dosage requirements. (see chart insert)
3. Mix with food to help ensure your dog receives a complete and proper dosing.
4. Make sure your dog eats all of the medicated food. You may need to moisten your dog's dry dog food to aid mixing.
5. The same dosage must be administered for three consecutive days. Do not stop treatment before the end of the three-day treatment period.