

**safe-guard**<sup>®</sup>  
(fenbendazole)

**This man is  
deworming  
his cattle.**

**No Chute. No Problem.**

Only SAFE-GUARD<sup>®</sup> (fenbendazole) pasture deworming options match the safety and efficacy of the drench and paste solutions producers have relied on for over 40 years.

**Let SAFE-GUARD do the work for you.**



**IMPORTANT SAFETY INFORMATION:** Do not use in beef calves less than 2 months old, dairy calves and veal calves. A withdrawal period has not been established for this product in pre-ruminating calves. Additionally, the following meat withdrawal and milk discard times apply: SAFE-GUARD<sup>®</sup> ENPROAL<sup>®</sup> Type C Medicated Block: Cattle must not be slaughtered for 11 days. For use in beef cattle only. SAFE-GUARD 20% Protein Type C Medicated Block: Cattle must not be slaughtered for 16 days. For use in beef cattle only. SAFE-GUARD Type A and other medicated feed products (pellets, cubes, free-choice mineral, or free-choice liquid): Cattle must not be slaughtered for 13 days. For dairy cattle, the milk discard time is 60 hours.

MAHCattle.com • 800-521-5767 • © 2023 Merck & Co., Inc., Rahway, NJ, USA and its affiliates. All rights reserved. US-SFG-230500004

